Historic, Archive Document

Do not assume content reflects current scientific knowledge, policies, or practices.



UNITED STATES DEPARTMENT OF AGRICULTURE PRODUCTION AND MARKETING ADMINISTRATION COTTON BRANCH BAKERSFIELD, CALIFORNIA

MOLUME XX, No. 9

MARCH 20, 1947

COTTON QUALITY REPORT FOR GINNINGS OF AMERICAN-EGYPTIAN 1946 CROP

TOTAL AMERICAN-EGYPTIAN (SXP) GINNINGS THIS SEASON CONTAINED A LARGER PROPORTION OF THE HIGHER GRADES AND LONGER STAPLES THAN LAST SEASON ACCORDING TO THE UNITED STATES DEPARTMENT OF AGRICULTURE. THE CURRENT CROP WAS ABOUT 53 PERCENT GRADES I AND 1-1/2, ALMOST 34 PERCENT GRADES 2 AND 2-1/2 AND ABOUT 13 PERCENT GRADE 3 AND LOWER. LAST SEASON THESE GRADE CATEGORIES WERE REPRESENTED BY PERCENTAGES OF APPROXIMATELY 42, 50 AND 8 RESPECTIVELY. STAPLES 1-1/2" AND LONGER ACCOUNTED FOR ALMOST 81 PERCENT OF THE 1946-47 GINNINGS COMPARED WITH ONLY ABOUT 67 PERCENT IN 1945-46.

ARIZONA-CALIFORNIA - THE GRADE DISTRIBUTION IN THIS SEASON'S GINNINGS CHANGED NOTICE-ABLY FROM A YEAR AGO. APPROXIMATELY 29 PERCENT OF THE CURRENT CROP WAS GRADES I AND 1-1/2, 48

PERCENT WAS GRADES 2 AND 2-1/2 AND ABOUT 23 PERCENT WAS GRADE 3 AND LOWER. LAST SEASON, PERCENTAGES FOR THESE GRADE CATEGORIES WERE APPROXIMATELY 24, 54 AND 12 RESPECTIVELY. THE PROPORTION OF STAPLES 1-1/2" AND LONGER INCREASED SHARPLY THIS SEASON, ALMOST 91 PERCENT AGAINST ONLY 58

PERCENT A YEAR EARLIER.

TEXAS, DISTRICT 6 - NEW MEXICO - THERE WAS LITTLE CHANGE IN THE GRADE DISTRIBUTION

FROM A YEAR AGO, HOWEVER, THE PROPORTION OF LONGER STAPLES DECREASED SHARPLY THIS SEASON.

APPROXIMATELY 82 PERCENT OF CURRENT GINNINGS WAS GRADES I AND 1-1/2 AND ABOUT 17 PERCENT WAS

GRADES 2 AND 2-1/2. LAST SEASON APPROXIMATELY 88 PERCENT WAS GRADES I AND 1-1/2 AND ABOUT 12

PERCENT WAS GRADES 2 AND 2-1/2. STAPLES 1-1/2" AND LONGER ACCOUNTED FOR ONLY ABOUT 68 PERCENT

OF THIS SEASON'S CROP COMPARED WITH ABOUT 89 PERCENT A YEAR AGO.

AMERICAN-EGYPTIAN (SXP) GINNINGS TOTALED 2,386 BALES ACCORDING TO THE BUREAU OF THE CENSUS.

THIS IS THE LAST GRADE AND STAPLE REPORT FOR ARIZONA-CALIFORNIA, TEXAS DISTRICT 6 - NEW MEXICO THIS SEASON.



1947	o u de es delle cui mange delle		1946 ** PcT.	-~~~	0.	7-1-6 3-1-6 1-6	0.	
MARCH 20, 1947	FAL	TOTAL	1946 PCT.	0 m	100.0	-17	100.0	
		AFTER JAN. 15	1945 PCT.	- 000- 000-700 004-7-04	100.0	200 200 200	100.0	IN COOPERATION WITH THE ARIZONA AGRICULTURAL EXPERIMENT STATION.
	TOTAL		1946** PCT.	るでででしょ 2 でいいっちゃ。 でいいの4がおい	100.0	06 0.0 4.0 4.0	100.0	
ESTIWATED GRADE AND STAPLE LENGTH OF AMERICAN-EGYPTIAN (SXP) COTTON GINNED			1945 PCT.	9.6 6.4 18.1 75.7 8.5	0.001	67.0	0.001	
	NEW MEXICO (B) & DIST. 6 OF TEXAS	AFTER TOTAL JAN. 15 CROP	1946** PCT.	40- 7044- 6-600	100.0	3.8 63.2 5.0	0.001	
			1945 PCT.	64- -40 vv-	100.0	89.0 89.4	100.0	
			1946** PCT.	3000	0.001	0.001	100.0	
			1945 PCT.	00	0.001	0.00	100.0	
	ARIZONA (A)-CALIFORNIA	TOTAL	1946** PCT.	000- 	100.0	89.5 1.6	100.0	COOPERATION
			1945 PCT.	4000004	100.0	21.04	100.0	(A) (B)
		AFTER JAN. 15	5 1946 * PCT.	2.2.2.2.2.2.2.2.2.2.2.2.2.2.2.2.2.2.2.	0.001 0.001	6/0 0-	0.001 0.001	
			1945 PCT.	80.000 80.400 80.4000	100.0	66.3	100.0	and the state of t
VOLUME XX, No. 9	en e	GRADE AND STAPLE LENGTH	GRADE	2-1/2 2-1/2 3-1/2 4-1/2 5 AND BELOW	ALL GRADES	LENGTH (INCHES) 1-9/32 AND SHORTER. 1-5/16 AND 1-11/32. 1-7/16 AND 1-15/32. 1-7/16 AND 1-17/32.	ALL LENGTHS	*LESS THAN 0.05 PERCENT